


## Direct measurement of the $7s^2S_{1/2} \rightarrow 7p^2P_{3/2}$ transition frequency in $^{226}\text{Ra}^+$

C. A. Holliman,\* M. Fan, A. Contractor, M. W. Straus, and A. M. Jayich  
 Department of Physics, University of California, Santa Barbara, California 93106, USA  
 and California Institute for Quantum Entanglement, Santa Barbara, California 93106, USA

 (Received 30 August 2020; accepted 8 October 2020; published 30 October 2020)

We report a direct measurement of the  $7s^2S_{1/2} \rightarrow 7p^2P_{3/2}$  electric dipole transition frequency in  $^{226}\text{Ra}^+$ . With a single laser-cooled radium ion we determine the transition frequency to be 785 722.11(3) GHz by directly driving the transition with frequency-doubled light and measuring the frequency of the undoubled light with an iodine reference. This measurement addresses a discrepancy of five combined standard deviations between previously reported values.

DOI: [10.1103/PhysRevA.102.042822](https://doi.org/10.1103/PhysRevA.102.042822)

Trapped radium ions are appealing for quantum information science [1], searches for new physics beyond the standard model [2,3], and precision timekeeping [4–6].  $\text{Ra}^+$  is a potential candidate for a transportable optical clock, as the transitions can all be addressed with direct diode lasers, and the ion’s laser cooling and fluorescence transition at 468 nm is far from the UV compared to most trapped ion optical clock candidates [7] (see Fig. 1). The narrow electric quadrupole transitions, the  $7s^2S_{1/2} \rightarrow 6d^2D_{5/2}$  transition at 728 nm and the  $7s^2S_{1/2} \rightarrow 6d^2D_{3/2}$  transition at 828 nm, may be used to study radium’s nuclear structure [8,9] or set bounds on sources of new physics [3].

There is a discrepancy between two previously reported values of the  $7s^2S_{1/2} \rightarrow 7p^2P_{3/2}$  (382-nm) transition frequency in  $^{226}\text{Ra}^+$ , 785 722.10(3) GHz [10] and 785 721.670(70) GHz [6], which is included in a compilation of radium spectroscopy data [11]. The former value was determined from a sum of two direct frequency measurements in  $^{226}\text{Ra}^+$  of the  $S_{1/2} \rightarrow D_{5/2}$  and  $D_{5/2} \rightarrow P_{3/2}$  transitions. The latter value was determined using the transition frequency calculated from a sum of direct measurements in  $^{214}\text{Ra}^+$  [6] and the isotope shift between  $^{214}\text{Ra}^+$  and  $^{226}\text{Ra}^+$  measured by Neu *et al.* [12]. In this work we help resolve this discrepancy with direct spectroscopy at 382 nm of the  $S_{1/2} \rightarrow P_{3/2}$  transition with a single laser-cooled radium ion. The frequency measurement is calibrated by absorption spectroscopy with known transitions in molecular iodine. The value measured in this work, 785 722.11(3) GHz, agrees with the value from the sum of direct measurements in  $^{226}\text{Ra}^+$  [10].

The iodine linear absorption spectrum is measured around 764 nm with light from a Ti:sapphire laser. Frequency-doubled light from the same laser drives the  $S_{1/2} \rightarrow P_{3/2}$  transition of  $^{226}\text{Ra}^+$  at 382 nm (see Fig. 2). From iodine reference lines near 764 nm we can determine the  $S_{1/2} \rightarrow P_{3/2}$  transition frequency. During both the radium and iodine spectroscopy the 764-nm laser’s frequency is recorded with a wavemeter (High Finesse WS-8) (see Fig. 3).

To measure the  $S_{1/2} \rightarrow P_{3/2}$  transition frequency we first scan the iodine spectrum near 764 nm, at half of the transition frequency reported in [10]. We scan over a 20-GHz range that includes two iodine reference lines, 325 and 326 [13]. This is followed by radium spectroscopy and then a second iodine scan to account for wavemeter drift. The radium spectroscopy uses state detection to determine if the population is in a “bright” state  $S_{1/2}, D_{3/2}$  or in the  $D_{5/2}$  “dark” state, by collecting  $\text{Ra}^+$  fluorescence at 468 nm onto a photomultiplier tube. The  $S_{1/2} \rightarrow P_{3/2}$  transition frequency is calculated from the difference between the radium 382-nm line center and twice the frequency of iodine line 326.

The pulse sequence for the  $S_{1/2} \rightarrow P_{3/2}$  frequency measurement is shown in Fig. 4. All light is linearly polarized to drive symmetric Zeeman transitions with the same amplitude, with a 3-G magnetic field along the trap’s axial direction. We Doppler cool the ion for 500  $\mu\text{s}$  before each pulse sequence. The initial state detection determines if the ion is cooled and if the population is in a bright state (SD1). If the population is not initialized in a bright state or if the ion is in a large orbit the data point is excluded. Population in the  $D_{3/2}$  state is then optically pumped for 100  $\mu\text{s}$  with light at 1079 nm to the ground state (P1). The  $S_{1/2} \rightarrow P_{3/2}$  spectroscopy transition is then driven with light at 382 nm for 500  $\mu\text{s}$  (P2). Decays from the  $P_{3/2}$  state have an 11% probability to be shelved in the  $D_{5/2}$  dark state [14]. We use these decays to determine the probability of driving the  $S_{1/2} \rightarrow P_{3/2}$  transition at a given laser frequency with a second state detection (SD2) to form a binomial distribution (see Fig. 5). Finally, any population that remains in the  $D_{5/2}$  state is reset to the bright states by optically pumping with 802-nm light (P3). The pulse sequence is repeated 32 000 times as the 382-nm laser is swept over the transition in a  $\sim 100$  MHz range. The  $S_{1/2} \rightarrow P_{3/2}$  spectroscopy is shown in Fig. 3, along with the corresponding iodine absorption reference spectrum.

The  $S_{1/2} \rightarrow P_{3/2}$  radium measurement data are binned by frequency and the shelving probability is fit to the exponential of a Lorentzian (see Fig. 5) to account for population depletion of the  $S_{1/2}$  state (see Appendix A). The full width at half maximum of 40(1) MHz gives a lower bound of 4.0(1) ns for the

\*cholliman3@gmail.com




FIG. 1. The  $\text{Ra}^+$  energy-level structure with the transitions driven in the measurement of the  $7s^2S_{1/2} \rightarrow 7p^2P_{3/2}$  transition frequency.

$P_{3/2}$  state lifetime, which is in agreement with the calculated value of 4.73 ns [15]. The 382-nm light is incident on the ion along the trap’s axial direction (see Fig. 2) to minimize micromotion broadening of the transition [16].

The closest iodine reference line to the  $^{226}\text{Ra}^+ S_{1/2} \rightarrow P_{3/2}$  transition is line 326 [13]. We calibrate the 326 line center frequency with a wavemeter to an absolute frequency using IodineSpec5 [17,18]. We fit the absorption dip of line 326 in the IodineSpec5 data to a Voigt function to determine its frequency-doubled line center, 785 734.265(3) GHz.

The frequency difference between the  $S_{1/2} \rightarrow P_{3/2}$  line center at 382 nm and twice the value of the iodine line 326 center frequency is 12.16(3) GHz (see Fig. 3). From this frequency difference we determine the  $S_{1/2} \rightarrow P_{3/2}$  transition frequency to be 785 722.11(3) GHz. The uncertainty is




FIG. 3. Iodine absorption (gray) and radium spectroscopy (red) data plotted in terms of the undoubled 764-nm laser frequency that we record on a wavemeter. Data are scaled and vertically offset for clarity. The iodine lines closest to the transition are lines 325 and 326 [13].

$\sigma_{\text{total}} = \sqrt{\sigma_{\text{Ra}^+}^2 + \sigma_{\text{I}_2}^2 + \sigma_{\text{spec}}^2 + \sigma_{\text{wm}}^2}$ , where  $\sigma_{\text{Ra}^+}$  (20 MHz) is half the full width at half maximum of the fitted  $\text{Ra}^+$  spectrum (see Fig. 5) to account for possible magnetic field shifts due to Zeeman levels,  $\sigma_{\text{I}_2}$  (10 MHz) is the iodine spectroscopy fitting uncertainty,  $\sigma_{\text{spec}}$  (3 MHz) is the IodineSpec5 line uncertainty, and  $\sigma_{\text{wm}}$  (20 MHz) is the wavemeter uncertainty.

This work measures a transition that was previously directly measured nearly a century ago [19] and addresses a discrepancy between the frequencies reported by Nuñez Portela *et al.* [6] and by Holliman *et al.* [10] (see Table I). This  $^{226}\text{Ra}^+$  frequency measurement also serves as a check for the  $S_{1/2} \rightarrow D_{5/2}$  (728 nm),  $S_{1/2} \rightarrow D_{3/2}$  (828 nm),  $D_{3/2} \rightarrow P_{3/2}$




FIG. 2. Diagram of the measurement setup. Light at 764 nm (red) from a Ti:sapphire laser is frequency doubled via second harmonic generation (SHG) to drive the  $S_{1/2} \rightarrow P_{3/2}$  transition in  $\text{Ra}^+$  at 382 nm (purple). An acousto-optic modulator (AOM) is used as a shutter for the 382-nm spectroscopy light. The 764-nm frequency is recorded on a wavemeter. A photodiode (PD) before the iodine cell compensates for laser power drifts. Other beams addressing the ions: Doppler cooling (teal), the  $D_{3/2} \rightarrow P_{1/2}$  repump (gray), and the  $D_{5/2} \rightarrow P_{3/2}$  cleanout (dark red).




FIG. 4. Pulse sequence used for the  $S_{1/2} \rightarrow P_{3/2}$  spectroscopy. Squiggly lines depict electric dipole allowed decays, straight lines with an arrow indicate optical pumping, and double arrows indicate optical cycling. The optical pumping (P) and state detection (SD) are described in the text.




FIG. 5. Spectroscopy of the  $S_{1/2} \rightarrow P_{3/2}$  transition at 382 nm. The Lorentzian fit that accounts for  $P_{3/2}$  state depletion gives a FWHM of 40(1) MHz and a reduced  $\chi^2$  of 1.06. Error bars are the most likely 68% confidence interval of a binomial distribution. The shelving probability is limited by branching fractions from the  $P_{3/2}$  state, as well as the probe duration, which was chosen to minimize power broadening while still realizing a reasonable signal-to-noise ratio.

(708 nm), and  $D_{5/2} \rightarrow P_{3/2}$  (802 nm) transition frequencies measured in [10]. The  $S_{1/2} \rightarrow P_{3/2}$  transition frequency can be calculated from the sum of the 728- and 802-nm frequencies, 785 722.10(3) GHz, or the 828- and 708-nm frequencies, 785 722.07(5) GHz. Both values derived from [10] agree with the reported direct measurement of the  $S_{1/2} \rightarrow P_{3/2}$  transition frequency. We verify the self-consistency of these sets of measurements in Appendix B.

We acknowledge use of the UC Santa Barbara physics department's Broadly-tunable Illumination Facility for Research, Outreach, Scholarship, and Training Ti:sapphire laser, which provided the 764- and 382-nm light used in this work. We also acknowledge support from NSF Grant No. PHY-1912665 and the UC Office of the President (Grant No. MRP-19-601445).

#### APPENDIX A: DEPLETION ANALYSIS

If a large proportion of the population is driven out of the  $S_{1/2}$  state during driving to the  $P_{3/2}$  state, the expected Lorentzian line shape of the spectroscopy distorts due to decays from the  $P_{3/2}$  state to the metastable  $D_{3/2}$  and  $D_{5/2}$  states. If we consider the  $S_{1/2}$  and  $P_{3/2}$  states as a two-level system and drive the  $S_{1/2} \rightarrow P_{3/2}$  transition for a time comparable to the  $P_{3/2}$  state lifetime or longer so that the system approaches equilibrium, there is a fraction of the population that remains in the excited state,  $\mathcal{R}_P$ . When we take into account decays to

TABLE I. Summary of reported  $S_{1/2} \rightarrow P_{3/2}$  transition frequencies in  $^{226}\text{Ra}^+$ . All values are offset from 785 722 000 MHz.

Transition	Ref. [19]	Ref. [6]	Ref. [10]	This work
$S_{1/2} \rightarrow P_{3/2}$	1000(4000)	-330(70) <sup>a</sup>	100(30) <sup>a</sup>	110(30)

<sup>a</sup>Frequency calculated from indirect measurements.

TABLE II. Summary of  $^{226}\text{Ra}^+$  transition frequency measurements. All units are GHz.

Transition	This work and [4,10]
$S_{1/2} \rightarrow P_{3/2}$	785 722.11(3)
$S_{1/2} \rightarrow P_{1/2}$	640 096.63(6)
$D_{3/2} \rightarrow P_{3/2}$	423 444.39(3)
$S_{1/2} \rightarrow D_{5/2}$	412 007.701(18)
$D_{5/2} \rightarrow P_{3/2}$	373 714.40(2)
$S_{1/2} \rightarrow D_{3/2}$	362 277.68(5)
$D_{3/2} \rightarrow P_{1/2}$	277 818.95(8) <sup>a</sup>

<sup>a</sup>Frequency calculated from a sum of direct measurements in  $^{226}\text{Ra}^+$ .

the metastable  $D_{3/2}$  and  $D_{5/2}$  states, the  $P_{3/2}$  state population  $\mathcal{P}_P$  is given by

$$\mathcal{P}_P = \mathcal{R}_P(1 - \mathcal{P}_D), \quad (\text{A1})$$

where  $\mathcal{P}_D$  is the total population in both the  $D_{3/2}$  and  $D_{5/2}$  states. We can approximate the decay into the  $D$  states as

$$\begin{aligned} \frac{d\mathcal{P}_D}{dt} &= \frac{\mathcal{P}_P(1-r)}{\tau} \\ &= \frac{\mathcal{R}_P(1-\mathcal{P}_D)(1-r)}{\tau}, \end{aligned} \quad (\text{A2})$$

where  $\tau$  is the lifetime of the  $P_{3/2}$  state and  $r$  is the branching fraction of the  $P_{3/2}$  state to the  $S_{1/2}$  state. Under the assumption that the population is always initialized to the  $S_{1/2}$  state, this gives

$$\mathcal{P}_D = 1 - \exp\left(\frac{-t\mathcal{R}_P(1-r)}{\tau}\right). \quad (\text{A3})$$

The final dark state population  $\mathcal{P}_{D_{s/2}}$  is determined from the branching fractions of the  $P_{3/2}$  state to the  $D_{5/2}$  state,  $s$ , and to the  $D_{3/2}$  state,  $t$ ,

$$\mathcal{P}_{D_{s/2}} = \frac{s}{s+t}\mathcal{P}_D, \quad (\text{A4})$$

which gives the fitting function

$$\mathcal{P}_{D_{s/2}} = \frac{s}{s+t} \left[ 1 - \exp\left(\frac{-a\Gamma}{(\omega - \omega_0)^2 + \Gamma^2}\right) + c \right], \quad (\text{A5})$$

where  $a$ ,  $\Gamma$ ,  $\omega$ , and  $c$  are fitting parameters and the  $s$  and  $t$  branching fractions are from [14].

#### APPENDIX B: RADIUM ION FREQUENCY SUMMARY

To verify that the measurements are self-consistent we perform an energy-level optimization with the reported  $S_{1/2} \rightarrow P_{3/2}$  transition frequency and other  $\text{Ra}^+$  transition frequencies reported in [4,10] (see Table II). If we fix the ground state at zero energy, there are seven transitions between four excited states which give three degrees of freedom. We use the energy-level optimization code package LOPT [20] to find the energy values that best agree with our reported frequencies. We calculate the residual sum of squares to be 0.14, which, being less than one, indicates that the measured frequencies are self-consistent.

- [1] D. Hucul, J. E. Christensen, E. R. Hudson, and W. C. Campbell, *Phys. Rev. Lett.* **119**, 100501 (2017).
- [2] N. Fortson, *Phys. Rev. Lett.* **70**, 2383 (1993).
- [3] J. C. Berengut, D. Budker, C. Delaunay, V. V. Flambaum, C. Frugiuele, E. Fuchs, C. Grojean, R. Harnik, R. Ozeri, G. Perez, and Y. Soreq, *Phys. Rev. Lett.* **120**, 091801 (2018).
- [4] M. Fan, C. A. Holliman, A. L. Wang, and A. M. Jayich, *Phys. Rev. Lett.* **122**, 223001 (2019).
- [5] B. K. Sahoo, R. G. E. Timmermans, B. P. Das, and D. Mukherjee, *Phys. Rev. A* **80**, 062506 (2009).
- [6] M. Nuñez Portela, E. A. Dijck, A. Mohanty, H. Bekker, J. E. van den Berg, G. S. Giri, S. Hoekstra, C. J. G. Onderwater, S. Schlessler, R. G. E. Timmermans, O. O. Versolato, L. Willmann, H. W. Wilschut, and K. Jungmann, *Appl. Phys. B* **114**, 173 (2014).
- [7] A. D. Ludlow, M. M. Boyd, J. Ye, E. Peik, and P. O. Schmidt, *Rev. Mod. Phys.* **87**, 637 (2015).
- [8] K. Heilig and A. Steudel, *At. Data Nucl. Data Tables* **14**, 613 (1974).
- [9] P.-G. Reinhard, W. Nazarewicz, and R. F. Garcia Ruiz, *Phys. Rev. C* **101**, 021301(R) (2020).
- [10] C. A. Holliman, M. Fan, and A. M. Jayich, *Phys. Rev. A* **100**, 062512 (2019).
- [11] U. Dammalapati, K. Jungmann, and L. Willmann, *J. Phys. Chem. Ref. Data* **45**, 013101 (2016).
- [12] W. Neu, R. Neugart, E. W. Otten, G. Passler, K. Wendt, B. Fricke, E. Arnold, H. J. Kluge, and G. Ulm, *Z. Phys. D* **11**, 105 (1989).
- [13] S. Gerstenkorn, J. Verges, and J. Chevillard, *Atlas du Spectre d'Absorption de la Molecule d'Iode: 11000–14000 cm to the Minus 1* (Laboratoire Aime-Cotton CNRS II, Paris, 1982).
- [14] M. Fan, C. A. Holliman, S. G. Porsev, M. S. Safronova, and A. M. Jayich, *Phys. Rev. A* **100**, 062504 (2019).
- [15] R. Pal, D. Jiang, M. S. Safronova, and U. I. Safronova, *Phys. Rev. A* **79**, 062505 (2009).
- [16] D. J. Berkeland, J. D. Miller, J. C. Bergquist, W. M. Itano, and D. J. Wineland, *J. Appl. Phys.* **83**, 5025 (1998).
- [17] H. Knöckel and E. Tiemann, *IodineSpec5* (Universität Hannover, Hannover, 2011).
- [18] H. Knöckel, B. Bodermann, and E. Tiemann, *Eur. Phys. J. D* **28**, 199 (2004).
- [19] E. Rasmussen, *Z. Phys.* **86**, 24 (1933).
- [20] A. Kramida, *Comput. Phys. Commun.* **182**, 419 (2011).